

In situ HCR v3.0 protocol for whole-mount chicken embryos (*Gallus gallus domesticus*)

This protocol has been optimized for embryos at stage HH 8–11, for other stages additional optimization may be required.

Technical support: support@moleculartechnologies.org

Preparation of fixed whole-mount chicken embryos

1. Collect chick embryos on 3M paper circles and place in a petri dish containing Ringer's solution.
2. Transfer embryos into a new petri dish with fresh Ringer's solution.
NOTE: This is to rinse away egg yolk before fixation.
3. Transfer into a petri dish containing 4% paraformaldehyde (PFA).
CAUTION: Use PFA with extreme care as it is a hazardous material.
NOTE: Use fresh PFA and cool to 4 °C before use to avoid increased autofluorescence.
4. Fix the samples at room temperature for 1 h.
5. Transfer embryos into a petri dish containing PBST.
6. Dissect the embryos off the filter paper. Remove the fixed vitelline membrane (opaque) and cut the squares around area pellucida, without leaving excess of extra-embryonic tissue.
7. Transfer embryos into a 2 mL eppendorf tube containing PBST.
8. Nutate for 5 mins with the tube positioned horizontally in a small ice bucket.
9. Wash embryos two additional times with 2 mL of PBST, each time nutating for 5 min on ice.
10. Dehydrate embryos with 2 × 5 min washes of 2 mL methanol (MeOH) on ice.
11. Store embryos at -20 °C overnight before use.
NOTE: Embryos can be stored for six months at -20 °C.
12. Transfer the required number of embryos for an experiment to a 2 mL eppendorf tube.
NOTE: Do not place more than 4 embryos in each 2 mL eppendorf tube.
13. Rehydrate with a series of graded 2 mL MeOH/PBST washes for 5 min each on ice:
 - (a) 75% MeOH / 25% PBST
 - (b) 50% MeOH / 50% PBST
 - (c) 25% MeOH / 75% PBST
 - (d) 100% PBST
 - (e) 100% PBST.
14. Treat embryos with 2 mL of 10 µg/mL proteinase K solution for 2 min (stage HH 8) or 2.5 min (stages HH 10–11) at room temperature.
NOTE: Proteinase K concentration and treatment time should be reoptimized for each batch of proteinase K, or for samples at a different developmental stage.
15. Postfix with 2 mL of 4% PFA for 20 min at room temperature.
CAUTION: Use PFA with extreme care as it is a hazardous material.
16. Wash embryos 2 × 5 min with 2 mL of PBST on ice.

17. Wash embryos with 2 mL of 50% PBST / 50% 5× SSCT for 5 min on ice.
18. Wash embryos with 2 mL of 5× SSCT for 5 min on ice.

Buffer recipes for sample preparation

4% Paraformaldehyde (PFA)

4% PFA
1× PBS

For 25 mL of solution

1 g of PFA powder
25 mL of 1× PBS
Heat solution at 50–60 °C to dissolve powder

PBST

1× PBS
0.1% Tween 20

For 50 mL of solution

5 mL of 10× PBS
500 μL of 10% Tween 20
Fill up to 50 mL with ultrapure H₂O

Proteinase K solution

10 μg/mL proteinase K

For 1 mL of solution

0.5 μL of 20 mg/mL proteinase K
Fill up to 1 mL with PBST

NOTE: Avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the embryos.

Multiplexed in situ HCR v3.0 protocol

Detection stage

1. For each sample, transfer 1-4 embryos to a 2 mL eppendorf tube.
NOTE: Do not place more than 4 embryos in each 2 mL eppendorf tube.
2. Incubate embryos in 1 mL of 30% probe hybridization buffer on ice for 5 min.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
3. Remove the buffer and pre-hybridize with 1 mL of 30% probe hybridization buffer for 30 min at 37 °C.
4. Prepare probe solution by adding 4 pmol of each probe mixture (odd & even: 2 μ L of 2 μ M stock per probe mixture) to 1 mL of 30% probe hybridization buffer at 37 °C.
NOTE: For dHCR imaging use 10 pmol of each probe to improve probe hybridization efficiency.
5. Remove the pre-hybridization solution and add the probe solution.
6. Incubate embryos overnight (12–16 h) at 37 °C.
7. Remove excess probes by washing with 4 \times 15 min with 1 mL of probe wash buffer at 37 °C:
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: Wash solutions should be pre-heated to 37 °C before use.
8. Wash samples 2 \times 5 min with 5 \times SSCT at room temperature.

Amplification stage

1. Pre-amplify embryos with 1 mL of amplification buffer for 5 min at room temperature.
2. Separately prepare 30 pmol of hairpin H1 and 30 pmol of hairpin H2 by snap cooling 10 μ L of 3 μ M stock (heat at 95 °C for 90 seconds and cool to room temperature in a dark drawer for 30 min).
NOTE: HCR hairpins H1 and H2 are provided in hairpin storage buffer ready for snap cooling. H1 and H2 should be snap cooled in separate tubes.
3. Prepare hairpin solution by adding all snap-cooled hairpins to 500 μ L of amplification buffer at room temperature.
4. Remove the pre-amplification solution and add the hairpin solution.
5. Incubate the embryos overnight (12–16 h) in the dark at room temperature.
NOTE: For dHCR imaging, amplify for 90-min to ensure single-molecule dots are diffraction-limited.
6. Remove excess hairpins by washing with 1 mL of 5 \times SSCT at room temperature:
 - (a) 2 \times 5 min
 - (b) 2 \times 30 min
 - (c) 1 \times 5 min

Sample mounting for microscopy

1. A chamber for mounting each embryo was made by aligning two stacks of double-sided tape (2 pieces per stack) 1 cm apart on a 25 mm × 75 mm glass slide.
2. Place an embryo between the tape stacks on the slide and remove as much solution as possible.
3. Align the embryo for dorsal imaging and carefully touch the slide with a kimwipe to further dry the area around the embryo.
4. Add two drops of SlowFade Gold antifade mountant on top of the embryo.
5. Place a 22 mm × 30 mm No. 1 coverslip on top of the stacks to close the chamber.
6. A Zeiss 710 NLO inverted confocal microscope equipped with an LD LCI Plan-Apochromat 25×/0.8 Imm Corr DIC objective was used in our lab to acquire chicken images.

Buffer recipes for in situ HCR

Probes, amplifiers, probe hybridization buffer, and probe wash buffer should be stored at -20 °C. Amplification buffer should be stored at 4 °C. Keep these reagents on ice at all times during probe and amplifier preparation. Make sure all solutions are well mixed before use.

30% probe hybridization buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin
 1× Denhardt's solution
 10% dextran sulfate

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 800 µL of 50× Denhardt's solution
 8 mL of 50% dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

30% probe wash buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 Fill up to 40 mL with ultrapure H₂O

Amplification buffer

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20
 10% dextran sulfate

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 8 mL of 50% dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 Fill up to 40 mL with ultrapure H₂O

50% dextran sulfate

50% dextran sulfate

For 40 mL of solution

20 g of dextran sulfate powder
 Fill up to 40 mL with ultrapure H₂O

Reagents and supplies

Paraformaldehyde (PFA) (Sigma Cat. # P6148)
Methanol (Mallinckrodt Chemicals Cat. # 3016-16)
Proteinase K, molecular biology grade (NEB Cat. # P8107S)
Formamide (Deionized) (Ambion Cat. # AM9342)
20× sodium chloride sodium citrate (SSC) (Life Technologies Cat. # 15557-044)
Heparin (Sigma Cat. # H3393)
50% Tween 20 (Life Technologies Cat. # 00-3005)
50× Denhardt's solution (Life Technologies Cat. # 750018)
Dextran sulfate, mol. wt. > 500,000 (Sigma Cat. # D6001)
25 mm × 75 mm glass slide (VWR Cat. # 48300-025)
22 mm × 30 mm No. 1 coverslip (VWR Cat. # 48393-026)
SlowFade Gold antifade mountant (Life Technologies Cat. # S36937)

Citation Notes

Molecular Technologies (moleculartechnologies.org) is a non-profit academic resource within the Beckman Institute at Caltech. For citation, please select from the list below as appropriate for your application:

- **In situ HCR v3.0**
Automatic background suppression for dramatically enhanced performance (signal-to-background and quantitative precision) and ease-of-use (no probe set optimization for new targets) (Choi *et al.*, 2018). v3.0 supports three quantitative modes: qHCR imaging – analog mRNA relative quantitation with subcellular resolution; qHCR flow cytometry – analog mRNA relative quantitation for high-throughput single-cell analysis; dHCR imaging – digital mRNA absolute quantitation. Protocols for v3.0 in diverse organisms are adapted from the zoo paper.
- **Quantitative HCR (qHCR) imaging**
mRNA relative quantitation with subcellular resolution in the anatomical context of whole-mount vertebrate embryos; read-out/read-in analysis framework (Trivedi *et al.*, 2018).
- **Zoo paper**
Protocols for multiplexed mRNA imaging in diverse sample types (Choi *et al.*, 2016):

bacteria in suspension	whole-mount zebrafish embryos and larvae
whole-mount fruit fly embryos	whole-mount chicken embryos
whole-mount worm larvae	whole-mount mouse embryos
whole-mount sea urchin embryos	FFPE human tissue sections
- **Single-molecule HCR imaging**
Single-molecule mRNA imaging in thick autofluorescent samples (e.g., 0.5 mm adult mouse brain sections) (Shah *et al.*, 2016).
- **Multiplexed quantitative HCR (qHCR) northern blots**
Simultaneous quantification of RNA target size and abundance with signal amplification for up to 5 target RNAs (Schwarzkopf & Pierce, 2016).
- **In situ HCR v2.0**
2nd generation in situ HCR technology (v2.0) using DNA probes and DNA HCR amplifiers: 10× increase in signal, 10× reduction in cost, dramatic increase in reagent durability (Choi *et al.*, 2014).
- **Shielded covalent (SC) probes**
Highly selective covalent capture of DNA and RNA targets, overcoming the longstanding tradeoff between selectivity and durable target capture (Vieregg *et al.*, 2013).
- **In situ HCR v1.0**
1st generation in situ HCR technology (v1.0) using RNA probes and RNA HCR amplifiers: multiplexed mRNA imaging in whole-mount vertebrate embryos with simultaneous signal amplification for up to 5 target mRNAs (Choi *et al.*, 2010).
- **Hybridization chain reaction (HCR) mechanism**
Isothermal enzyme-free molecular signal amplification (Dirks & Pierce, 2004).

We are happy to provide advice and technical support: support@moleculartechnologies.org.

– The Molecular Technologies Team

References

- Choi, H. M. T., Chang, J. Y., Trinh, L. A., Padilla, J. E., Fraser, S. E., & Pierce, N. A. (2010). Programmable in situ amplification for multiplexed imaging of mRNA expression. *Nat Biotechnol*, **28**(11), 1208–12. ([pdf](#), [supp info](#), [supp movie 1](#), [supp movie 2](#))
- Choi, H. M. T., Beck, V. A., & Pierce, N. A. (2014). Next-generation in situ hybridization chain reaction: higher gain, lower cost, greater durability. *ACS Nano*, **8**(5), 4284–4294. ([pdf](#), [supp info](#), [supp movie](#))
- Choi, H. M. T., Calvert, C. R., Husain, N., Huss, D., Barsi, J. C., Deverman, B. E., Hunter, R. C., Kato, M., Lee, S. M., Abelin, A. C. T., Rosenthal, A. Z., Akbari, O. S., Li, Y., Hay, B. A., Sternberg, P. W., Patterson, P. H., Davidson, E. H., Mazmanian, S. K., Prober, D. A., van de Rijn, M., Leadbetter, J. R., Newman, D. K., Readhead, C., Bronner, M. E., Wold, B., Lansford, R., Sauka-Spengler, T., Fraser, S. E., & Pierce, N. A. (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, **143**, 3632–3637. ([pdf](#), [supp info](#), [fruit fly movie](#), [sea urchin movie](#), [zebrafish movie](#), [chicken movie](#), [mouse movie](#))
- Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., & Pierce, N. A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development*, **145**, doi:10.1242/dev.165753. ([pdf](#), [supp info](#), [Dot Analysis 1.0 package](#))
- Dirks, R. M., & Pierce, N. A. (2004). Triggered amplification by hybridization chain reaction. *Proc Natl Acad Sci USA*, **101**(43), 15275–15278. ([pdf](#))
- Schwarzkopf, M., & Pierce, N. A. (2016). Multiplexed miRNA northern blots via hybridization chain reaction. *Nucleic Acids Res*, **44**(15), e129. ([pdf](#), [supp info](#))
- Shah, S., Lubeck, E., Schwarzkopf, M., He, T.-F., Greenbaum, A., Sohn, C. H., Lignell, A., Choi, H. M. T., Gradinaru, V., Pierce, N. A., & Cai, L. (2016). Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. *Development*, **143**, 2862–2867. ([pdf](#), [supp info](#), [supp movie 1](#), [supp movie 2](#), [supp movie 3](#))
- Trivedi, V., Choi, H. M. T., Fraser, S. E., & Pierce, N. A. (2018). Multidimensional quantitative analysis of mRNA expression within intact vertebrate embryos. *Development*, **145**, doi:10.1242/dev.156869. ([pdf](#), [supp info](#), [Read-out/Read-in 1.0 package](#))
- Vieregg, J. R., Nelson, H. M., Stoltz, B. M., & Pierce, N. A. (2013). Selective nucleic acid capture with shielded covalent probes. *J Am Chem Soc*, **135**(26), 9691–9699. ([pdf](#), [supp info](#))