

In situ HCR v3.0 protocol for tissue section on slide

This protocol has not been validated for all sample types yet and should only be used as a template.
Technical support: support@moleculartechnologies.org

Sample pre-treatment

Samples should be prepared in the same manner as for a traditional in situ hybridization, up to the probe hybridization step. This may include permeabilization and protease digestion. Then proceed with the protocol described below.

Multiplexed in situ HCR v3.0 protocol

Detection stage

1. Pre-warm a humidified chamber to 37 °C.
2. Dry slide by blotting edges on a Kimwipe.
3. Add 200 μL of 30% probe hybridization buffer on top of the tissue sample.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
4. Pre-hybridize for 10 min inside the humidified chamber.
5. Prepare probe solution by adding 0.4 pmol of each probe mixture (odd & even: 0.2 μL of 2 μM stock per probe mixture) to 100 μL of probe hybridization buffer at 37 °C.
NOTE: For dHCR imaging, use higher concentration of probe to improve probe hybridization efficiency.
6. Remove the pre-hybridization solution and drain excess buffer on slide by blotting edges on a Kimwipe.
7. Add 50–100 μL of the probe solution on top of the tissue sample.
NOTE: Amount of probe solution depends on the size of the coverslip.
8. Place a coverslip on the tissue sample and incubate overnight (12–16 h) in the 37 °C humidified chamber.
9. Immerse slide in probe wash buffer at 37 °C to float off coverslip.
CAUTION: probe wash buffer contains formamide, a hazardous material.
10. Remove excess probes by incubating slide at 37 °C in:
 - (a) 75% of 30% probe wash buffer / 25% 5 \times SSCT for 15 min
 - (b) 50% of 30% probe wash buffer / 50% 5 \times SSCT for 15 min
 - (c) 25% of 30% probe wash buffer / 75% 5 \times SSCT for 15 min
 - (d) 100% 5 \times SSCT for 15 min*NOTE: Wash solutions should be pre-heated to 37 °C before use.*
11. Immerse slide in 5 \times SSCT for 5 min at room temperature.

Amplification stage

1. Dry slide by blotting edges on a Kimwipe.
2. Add 200 μL of amplification buffer on top of the tissue sample and pre-amplify in a humidified chamber for 30 min at room temperature.
3. Separately prepare 6 pmol of hairpin H1 and 6 pmol of hairpin H2 by snap cooling 2 μL of 3 μM stock (heat at 95 $^{\circ}\text{C}$ for 90 seconds and cool to room temperature in a dark drawer for 30 min).
NOTE: HCR hairpins H1 and H2 are provided in hairpin storage buffer ready for snap cooling. H1 and H2 should be snap cooled in separate tubes.
4. Prepare hairpin solution by adding all snap-cooled hairpins to 100 μL of amplification buffer at room temperature.
5. Remove the pre-amplification solution and drain excess buffer on slide by blotting edges on a Kimwipe.
6. Add 50–100 μL of the hairpin solution on top of the tissue sample.
NOTE: Amount of hairpin solution depends on the size of the coverslip.
7. Place a coverslip on the tissue sample and incubate overnight (12–16 h) in a dark humidified chamber at room temperature.
NOTE: For dHCR imaging, amplify for a shorter period of time to ensure single-molecule dots are diffraction-limited.
8. Immerse slide in $5 \times$ SSCT at room temperature to float off coverslip.
9. Remove excess hairpins by incubating slide in $5 \times$ SSCT at room temperature for:
 - (a) 2×30 min
 - (b) 1×5 min

Sample mounting for microscopy

1. Dry slide by blotting edges on a Kimwipe.
2. Add 50 μL of SlowFade Gold antifade mountant with DAPI on top of tissue section.
3. Place a coverslip on top for imaging.
4. A Zeiss Axio Observer Z1 fluorescent inverted microscope equipped with an Plan-Apochromat 63 \times /1.4 Oil DIC M27 objective was used in our lab to acquire tissue section images. Samples were illuminated with X-Cite Series 120 (EXFO Photonics) and images were collected with an ImagEM-1K EM-CCD digital camera (Hamamatsu).

Buffer recipes for in situ HCR

Probes, amplifiers, probe hybridization buffer, and probe wash buffer should be stored at -20 °C. Amplification buffer should be stored at 4 °C. Keep these reagents on ice at all times during probe and amplifier preparation. Make sure all solutions are well mixed before use.

30% probe hybridization buffer

30% formamide
5× sodium chloride sodium citrate (SSC)
9 mM citric acid (pH 6.0)
0.1% Tween 20
50 µg/mL heparin
1× Denhardt's solution
10% dextran sulfate

For 40 mL of solution

12 mL formamide
10 mL of 20× SSC
360 µL 1 M citric acid, pH 6.0
400 µL of 10% Tween 20
200 µL of 10 mg/mL heparin
800 µL of 50× Denhardt's solution
8 mL of 50% dextran sulfate
Fill up to 40 mL with ultrapure H₂O

30% probe wash buffer

30% formamide
5× sodium chloride sodium citrate (SSC)
9 mM citric acid (pH 6.0)
0.1% Tween 20
50 µg/mL heparin

For 40 mL of solution

12 mL formamide
10 mL of 20× SSC
360 µL 1 M citric acid, pH 6.0
400 µL of 10% Tween 20
200 µL of 10 mg/mL heparin
Fill up to 40 mL with ultrapure H₂O

Amplification buffer

5× sodium chloride sodium citrate (SSC)
0.1% Tween 20
10% dextran sulfate

For 40 mL of solution

10 mL of 20× SSC
400 µL of 10% Tween 20
8 mL of 50% dextran sulfate
Fill up to 40 mL with ultrapure H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
400 µL of 10% Tween 20
Fill up to 40 mL with ultrapure H₂O

50% dextran sulfate

50% dextran sulfate

For 40 mL of solution

20 g of dextran sulfate powder
Fill up to 40 mL with ultrapure H₂O

Reagents and supplies

Formamide (Deionized) (Ambion Cat. # AM9342)

20× sodium chloride sodium citrate (SSC) (Life Technologies Cat. # 15557-044)

Heparin (Sigma Cat. # H3393)

50× Denhardt's solution (Life Technologies Cat. # 750018)

Dextran sulfate, mol. wt. > 500,000 (Sigma Cat. # D6001)

10% Tween 20 Solution (Bio-Rad Cat. # 161-0781)

SlowFade Gold antifade mountant with DAPI (Life Technologies Cat. # S36938)

Citation Notes

Molecular Technologies (moleculartechnologies.org) is a non-profit academic resource within the Beckman Institute at Caltech. For citation, please select from the list below as appropriate for your application:

- **In situ HCR v3.0**
Automatic background suppression for dramatically enhanced performance (signal-to-background and quantitative precision) and ease-of-use (no probe set optimization for new targets) (Choi *et al.*, 2018). v3.0 supports three quantitative modes: qHCR imaging – analog mRNA relative quantitation with subcellular resolution; qHCR flow cytometry – analog mRNA relative quantitation for high-throughput single-cell analysis; dHCR imaging – digital mRNA absolute quantitation. Protocols for v3.0 in diverse organisms are adapted from the zoo paper.
- **Quantitative HCR (qHCR) imaging**
mRNA relative quantitation with subcellular resolution in the anatomical context of whole-mount vertebrate embryos; read-out/read-in analysis framework (Trivedi *et al.*, 2018).
- **Zoo paper**
Protocols for multiplexed mRNA imaging in diverse sample types (Choi *et al.*, 2016):

bacteria in suspension	whole-mount zebrafish embryos and larvae
whole-mount fruit fly embryos	whole-mount chicken embryos
whole-mount worm larvae	whole-mount mouse embryos
whole-mount sea urchin embryos	FFPE human tissue sections
- **Single-molecule HCR imaging**
Single-molecule mRNA imaging in thick autofluorescent samples (e.g., 0.5 mm adult mouse brain sections) (Shah *et al.*, 2016).
- **Multiplexed quantitative HCR (qHCR) northern blots**
Simultaneous quantification of RNA target size and abundance with signal amplification for up to 5 target RNAs (Schwarzkopf & Pierce, 2016).
- **In situ HCR v2.0**
2nd generation in situ HCR technology (v2.0) using DNA probes and DNA HCR amplifiers: 10× increase in signal, 10× reduction in cost, dramatic increase in reagent durability (Choi *et al.*, 2014).
- **Shielded covalent (SC) probes**
Highly selective covalent capture of DNA and RNA targets, overcoming the longstanding tradeoff between selectivity and durable target capture (Vieregg *et al.*, 2013).
- **In situ HCR v1.0**
1st generation in situ HCR technology (v1.0) using RNA probes and RNA HCR amplifiers: multiplexed mRNA imaging in whole-mount vertebrate embryos with simultaneous signal amplification for up to 5 target mRNAs (Choi *et al.*, 2010).
- **Hybridization chain reaction (HCR) mechanism**
Isothermal enzyme-free molecular signal amplification (Dirks & Pierce, 2004).

We are happy to provide advice and technical support: support@moleculartechnologies.org.

– The Molecular Technologies Team

References

- Choi, H. M. T., Chang, J. Y., Trinh, L. A., Padilla, J. E., Fraser, S. E., & Pierce, N. A. (2010). Programmable in situ amplification for multiplexed imaging of mRNA expression. *Nat Biotechnol*, **28**(11), 1208–12. ([pdf](#), [supp info](#), [supp movie 1](#), [supp movie 2](#))
- Choi, H. M. T., Beck, V. A., & Pierce, N. A. (2014). Next-generation in situ hybridization chain reaction: higher gain, lower cost, greater durability. *ACS Nano*, **8**(5), 4284–4294. ([pdf](#), [supp info](#), [supp movie](#))
- Choi, H. M. T., Calvert, C. R., Husain, N., Huss, D., Barsi, J. C., Deverman, B. E., Hunter, R. C., Kato, M., Lee, S. M., Abelin, A. C. T., Rosenthal, A. Z., Akbari, O. S., Li, Y., Hay, B. A., Sternberg, P. W., Patterson, P. H., Davidson, E. H., Mazmanian, S. K., Prober, D. A., van de Rijn, M., Leadbetter, J. R., Newman, D. K., Readhead, C., Bronner, M. E., Wold, B., Lansford, R., Sauka-Spengler, T., Fraser, S. E., & Pierce, N. A. (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, **143**, 3632–3637. ([pdf](#), [supp info](#), [fruit fly movie](#), [sea urchin movie](#), [zebrafish movie](#), [chicken movie](#), [mouse movie](#))
- Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., & Pierce, N. A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development*, **145**, doi:10.1242/dev.165753. ([pdf](#), [supp info](#), [Dot Analysis 1.0 package](#))
- Dirks, R. M., & Pierce, N. A. (2004). Triggered amplification by hybridization chain reaction. *Proc Natl Acad Sci USA*, **101**(43), 15275–15278. ([pdf](#))
- Schwarzkopf, M., & Pierce, N. A. (2016). Multiplexed miRNA northern blots via hybridization chain reaction. *Nucleic Acids Res*, **44**(15), e129. ([pdf](#), [supp info](#))
- Shah, S., Lubeck, E., Schwarzkopf, M., He, T.-F., Greenbaum, A., Sohn, C. H., Lignell, A., Choi, H. M. T., Gradi-naru, V., Pierce, N. A., & Cai, L. (2016). Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. *Development*, **143**, 2862–2867. ([pdf](#), [supp info](#), [supp movie 1](#), [supp movie 2](#), [supp movie 3](#))
- Trivedi, V., Choi, H. M. T., Fraser, S. E., & Pierce, N. A. (2018). Multidimensional quantitative analysis of mRNA expression within intact vertebrate embryos. *Development*, **145**, doi:10.1242/dev.156869. ([pdf](#), [supp info](#), [Read-out/Read-in 1.0 package](#))
- Vieregg, J. R., Nelson, H. M., Stoltz, B. M., & Pierce, N. A. (2013). Selective nucleic acid capture with shielded covalent probes. *J Am Chem Soc*, **135**(26), 9691–9699. ([pdf](#), [supp info](#))